

Tel: +1(800)421-4149, Fax: +1(510)824-1490, support@biogenex.com

# DATA SHEET Hsa-miR-9fluoresceinatedoligo probe

Catalog No. Description

**HM009-100** One vial of 0.650 ml of probe in hybridization buffer

## Analyte Specific Reagent. Analytical and performance characteristics are not established.

Doc. No. 932-HM009-100 Rev: C Date of release: 10-Aug-2020

#### **Description**

The Hsa-miR-9probe has been designed from mature human miR-9sequence. This fluoresceinated probe is provided in a hybridization buffer for localization of miRNA in FFPE tissue by *In Situ* hybridization.

## **Specifications**

The Hsa-miR-9identifies mature miR-9sequences in formalin-fixed, paraffin-embedded human tissues and/orfreshly prepared frozen tissues by *in situ* hybridization. This probe does not react with normal human mRNA or nuclear DNA present in tissues.

#### **Storage and Handling**

Store the reagent at 2-8 °C. Do not freeze. Do not use the reagent after expiration date on vial. The reagent must be brought to room temperature before use. (Important! The presence of precipitates induces background staining).

## **Precautions:**

For professional use. The probe contains formamide. Formamide is classified as a teratogen. Pregnant workers should keep exposure to a minimum. Avoid inhalation, ingestion, and contact with unprotected skin. If skin contact occurs, wash thoroughly with soap and water. For more information, refer to the Material Safety Data Sheet, which is available upon request.

#### **Quality Control**

Each lot of this micro RNA probe is tested by *In Situ* hybridization for Quality Control purposes. Refer to the BioGenex Quality Control Testing Conditions table for additional information.

#### References

- 1. Nasser MW. et al. Down-regulation of Micro-RNA-1 (miR-1) in Lung Cancer. THE JOURNAL OF BIOLOGICAL CHEMISTRY 283 (48), 33394–33405 (2008).
- 2. Datta J. et al. Methylation Mediated Silencing of MicroRNA-1 Gene and Its Role in Hepatocellular Carcinogenesis. *Cancer Res* 68, 5049-5058 (2008).
- 3. Kloosterman WP.et al.*in situ* detection of miRNAs in animal embryos using LNA-modified oligonucleotide probes. Nature Methods, 3, 27 29 (2006).
- 4. Wheeler G. et al. *In situ* detection of animal and plant microRNAs. *DNA Cell Biol*, **26**, 251–255 (2007).
- 5. Nuovo GJ. In situ detection of precursor and mature microRNAs in paraffin embedded, formalinfixed tissues and cell preparations. Methods 44(1),39–46 (2008).
- 6. Song R. et al. In situ hybridization detection of microRNAs. Methods Mol Biol. 629, 287-94 (2010).

#### **BioGenex Quality Control Testing Conditions**

Parameter	Conditions used
Control Tissue	STOMACH Ca., COLON Ca. (FB-HM0090)
Tissue Type	Formalin-fixed, paraffin-embedded cancer tissues