

**DATA SHEET**  
**Hsa-miR-150 fluoresceinated oligo probe**

<b>Catalog No.</b>	<b>Description</b>
<b>HM150-100</b>	One vial of 0.650 ml of probe in hybridization buffer

**Analyte Specific Reagent. Analytical and performance characteristics are not established.**

Doc. No. 932-HM150-100 Rev C

Date of release: 12-Aug-2020

**Description**

The Hsa-miR-150 probe has been designed from mature human miR-150 sequence. This fluoresceinated probe is provided in a hybridization buffer for localization of miRNA in FFPE tissue by *In Situ* hybridization.

**Specifications**

The Hsa- miR-150 identifies mature miR-150 sequences in formalin-fixed, paraffin-embedded human tissues and freshly prepared frozen tissues by *in situ* hybridization. This probe does not react with normal human mRNA or nuclear DNA present in tissues.

**Storage and Handling**

Store the reagent at 2-8 °C. Do not freeze. Do not use after expiration date on vial. The reagent must be brought to room temperature before use. (Important! The presence of precipitates induces background staining).

**Precautions:**

For professional use. The probe contains formamide. Formamide is classified as a teratogen. Pregnant workers should keep exposure to a minimum. Avoid inhalation, ingestion, and contact with unprotected skin. If skin contact occurs, wash thoroughly with soap and water. For more information, refer to the Material Safety Data Sheet, which is available upon request.

**Quality Control**

Each lot of this micro RNA probe is tested by *In Situ* hybridization for Quality Control purposes. Refer to the BioGenex Quality Control Testing Conditions table for additional information.

**References**

1. Lin YC et al. c-Myb is an evolutionary conserved miR-150 target and miR-150/c-Myb interaction is important for embryonic development. *Mol Biol Evol* 25 (10): 2189-2198 (2008).
2. Kloosterman WP et al. *in situ* detection of miRNAs in animal embryos using LNA-modified oligonucleotide probes. *Nature Methods*, 3, 27 – 29 (2006).
3. Wheeler G. et al. *In situ* detection of animal and plant microRNAs. *DNA Cell Biol*, 26, 251–255 (2007).
4. Nuovo GJ. In situ detection of precursor and mature microRNAs in paraffin embedded, formalin fixed tissues and cell preparations. *Methods* 44(1), 39–46 (2008).
5. Song R. et al. *In situ* hybridization detection of microRNAs. *Methods Mol Biol*. 629, 287-94 (2010)

**BioGenex Quality Control Testing Conditions**

<b>Parameter</b>	<b>Conditions used</b>
Control Tissue	LYMPH NODE TESTIS (FB-HM150).
Tissue Type	Formalin-fixed, paraffin-embedded cancer tissues