

48810 Kato Road, Suite 100E & 200E Fremont, CA 94538

support@biogenex.com

Tel: +1 (800) 421-4149, Fax: +1 (510) 824-1490,

DATA SHEET Hsa-miR-218 fluoresceinated oligo probe

Catalog No. HM218-100

Description

One vial of 0.650 ml of probe in hybridization buffer

Analyte Specific Reagent. Analytical and performance characteristics are not established.

Doc. No. 932-HM218-100 Rev:C

Date of release: 14-Aug-2020

Description

The Hsa-miR-218 probe has been designed from mature human miR-218 sequence. This fluoresceinated probe is provided in a hybridization buffer for localization of miRNA in FFPE tissue by *In Situ* hybridization.

Specifications

The Hsa-miR-218 identifies mature miR-218 sequences in formalin-fixed, paraffin-embedded human tissues and/or freshly prepared frozen tissues by *in situ* hybridization. This probe does not react with normal human mRNA or nuclear DNA present in tissues.

Storage and Handling

Store the reagent at 2-8 °C. Do not freeze. Do not use the reagent after expiration date on vial. The reagent must be brought to room temperature before use. (Important! The presence of precipitates induces background staining).

Precautions:

For professional use the probe contains formamide. Formamide is classified as a teratogen. Pregnant workers should keep exposure to a minimum. Avoid inhalation, ingestion, and contact with unprotected skin. If skin contact occurs, wash thoroughly with soap and water. For more information, refer to the Material Safety Data Sheet, which is available upon request.

Quality Control

Each lot of this micro RNA probe is tested by *In-Situ* hybridization for Quality Control purposes. Refer to the BioGenex Quality Control Testing Conditions table for additional information.

References

- 1. Nasser MW. et al. Down-regulation of Micro-RNA-1 (miR-1) in Lung Cancer. THE JOURNAL OF BIOLOGICAL CHEMISTRY 283 (48), 33394–33405 (2008).
- 2. Datta J. et al. Methylation Mediated Silencing of MicroRNA-1 Gene and Its Role in Hepatocellular Carcinogenesis. *Cancer Res* 68, 5049-5058 (2008).
- 3. Kloosterman WP. et al. *in situ* detection of miRNAs in animal embryos using LNA-modified oligonucleotide probes. Nature Methods, 3, 27 29 (2006).
- 4. Wheeler G. et al. *In situ* detection of animal and plant microRNAs. *DNA Cell Biol*, **26**, 251–255 (2007).
- 5. Nuovo GJ. In situ detection of precursor and mature microRNAs in paraffin embedded, formalin fixed tissues and cell preparations. Methods 44(1),39–46 (2008).
- 6. Song R. et al. In situ hybridization detection of microRNAs. Methods Mol Biol. 629, 287-94 (2010).

BioGenex Quality Control Testing Conditions

Parameter	Conditions used
Control Tissue	NORMAL CERVIX/ BREAST CARCINOMA (FB-HM218).
Tissue Type	Formalin-fixed, paraffin-embedded cancer tissues